



# Coral Bleaching Early Warning Network

## Current Conditions Report #20190920



Updated September 20, 2019

**Summary:** Based on climate predictions, current conditions, and field observations, the threat for mass coral bleaching within the FKNMS remains **LOW**.

### NOAA Coral Reef Watch Current and 60% Probability Coral Bleaching Alert Outlook September 18, 2019 (experimental)




Figure 1. NOAA's 5 km Experimental Current and 60% Probability Coral Bleaching Alert Outlook Areas through December 2019. Updated September 18, 2019. [http://coralreefwatch.noaa.gov/vs/gauges/florida\\_keys.php](http://coralreefwatch.noaa.gov/vs/gauges/florida_keys.php)




Figure 2. NOAA's Experimental 5km Coral Bleaching HotSpot Map for Florida September 18, 2019. [https://coralreefwatch.noaa.gov/vs/gauges/florida\\_keys.php](https://coralreefwatch.noaa.gov/vs/gauges/florida_keys.php)




Figure 3. NOAA's Experimental 5km Degree Heating Weeks Map for Florida September 18, 2019. [https://coralreefwatch.noaa.gov/vs/gauges/florida\\_keys.php](https://coralreefwatch.noaa.gov/vs/gauges/florida_keys.php)

### Weather and Sea Temperatures

According to the newly released NOAA Coral Reef Watch (CRW) experimental 5 kilometer (km) Satellite Current and 60% Probability Coral Bleaching Alert Area, all areas of the Florida Keys National Marine Sanctuary are under a bleaching Watch, however, the coral bleaching outlook conditions are currently not favorable for a mass bleaching event (Fig. 1).

Recent remote sensing analysis by NOAA's CRW program indicates that all of the Florida Keys region is currently experiencing minimal thermal stress. NOAA's new experimental 5 km Coral Bleaching HotSpot Map (Fig. 2), which illustrates current sea surface temperatures compared to the average temperature for the warmest month, shows slightly elevated temperatures for the Florida Keys. Similarly, NOAA's experimental 5 km Degree Heating Weeks (DHW) map, which illustrates how much heat stress has built up over the past 12 weeks (Fig.3), indicates limited temperature stress is currently evident in the Florida Keys region. NOAA's Integrated Coral Observing Network (ICON) and Pacific Marine Environmental Laboratory (PMEL) monitoring stations, which provide near real time *in-situ* sea temperature and/or wind data throughout the Florida Keys reefs, as well as Mote Marine Laboratory (MML) *in-situ* temperature data confirm that temperatures remain at or just below 30°C over the past week (Fig.4) likely due in part to moderate to high wind conditions (Fig. 5). Mote Marine Laboratory will continue to monitor the NOAA HotSpot maps, DHW maps, and ICON sea temperature data from NOAA monitoring stations on a weekly basis for the remainder of the bleaching season.




Figure 4. *in-situ* sea temperature from NOAA/ICON and MML monitoring stations (Sept. 1-20, 2019).




Figure 5. Wind speed data from NOAA/ICON monitoring stations (Sept. 1-20, 2019).



### Current Coral Conditions

A total of 51 BleachWatch Observer reports were received during the last 2 weeks (Fig. 6), with 39 reports indicating isolated colonies exhibiting signs of paling or partial bleaching (Fig. 7) and 9 reports of isolated bleaching (Fig. 8) in the Dry Tortugas. The remaining 3 reports indicated that no significant signs of coral bleaching were observed. At those sites where paling/partial bleaching was noted, the overall percentage of corals exhibiting signs of thermal stress was mostly 11-30%, however several inshore/shallow reef sites noted up to 75% of corals affected. The majority of paling/partial bleaching observations consisted of isolated colonies of Encrusting/Mound/Boulder corals (*Siderastrea siderea*, *Siderastrea radians*, *Montastraea cavernosa*, *Orbicella faveolata*, *Orbicella annularis*, *Porities ssp.* and *Dichocoenia stokesii*), Brain corals (*Colpohyllia natans*, *Pseudodiploria strigosa*, *Pseudodiploria clivosa* and *Diploria labyrinthiformis*) and Leaf/Plate/Sheet corals (*Agaricia spp.*) Other observations included paling of *Palythoa spp.*, Fire Coral, and Gorgonians as well as numerous reports of coral disease.



Figure 7. Paling *C. natans* at a nearshore reef off Cudjoe Key on 9/18/19.




Figure 8. Bleached *A. agaricites* Wonderland on 9/18/19.

These isolated observations of paling and partial bleaching do not necessarily indicate that the onset of a mass bleaching event is currently underway; however, continued field observations are needed as more widespread coral bleaching could potentially develop if environmental conditions continue to be favorable.

### BleachWatch Reports for Sept. 4-19, 2019




Figure 6. Overview of BleachWatch observer reports submitted from Sept. 4-19, 2019

**For more information about the BleachWatch program,  
or to submit a bleaching observation, contact:**



Cory Walter  
Mote Marine Laboratory  
24244 Overseas Highway  
Summerland Key, FL 33042  
(305) 395-8730

<http://www.mote.org/bleachwatch>

**FUNDING THANKS TO....**

